

## Effect of previous water conditions on vine response to rewatering

M. GÓMEZ-DEL-CAMPO, P. BAEZA, C. RUIZ, V. SOTÉS and J. R. LISSARRAGUE

Departamento de Producción Vegetal: Fitotecnia, Universidad Politécnica de Madrid, Madrid, Spain

### Summary

**A comparative study of stomatal responses to rewatering was conducted on grapevines previously subjected to conditions of water stress and no water stress. Two cultivars were grown in 35 L lysimeters: 'Airén', from the dry zone of La Mancha, and 'Chardonnay', from the humid zone of Burgundy. The day after rewatering no significant differences in soil water content were found between water treatments or cultivars. However, predawn leaf water potential was significantly higher under non-stress than under stressed conditions, differences between cultivars were also found. Water consumption during the 5 d after rewatering was directly related to vine leaf area development. Vines grown under water stress conditions showed more uniform stomatal behaviour after rewatering. One day after rewatering rates of leaf conductance, transpiration and net photosynthesis were significantly higher in non-stressed vines, while 3 d after rewatering leaf conductance and transpiration were significantly higher in vines previously subjected to water stress. Net photosynthesis was significantly higher in water-stressed vines 5 d after rewatering. No differences were found between cultivars with regard to leaf conductance. The differences in the relationship between net photosynthesis and leaf conductance between stressed and non-stressed vines before rewatering were not found after rewatering. The only permanent adaptation mechanism to water stress was a lowering of leaf area development, which allowed water-stressed vines to consume less water to maintain a higher water availability and high or constant stomatal conductance.**

**Abbreviations:**  $\Psi_{PD}$  - predawn leaf water potential; A - net photosynthesis;  $g_l$  - leaf conductance; E - transpiration; NS - no water stress; S - water stress.

**Key words:** *Vitis vinifera* L., cv. Airén, cv. Chardonnay, predawn leaf water potential, photosynthesis, leaf conductance, transpiration.

### Introduction

Under mediterranean climatic conditions, soil moisture content slowly decreases from the beginning of spring to summer. In some years intermittent rainfall is common throughout summer. Plant growth and survival after these rains depend on the plant's capacity to use available water efficiently and its response to the following drought period.

During the season it is difficult to determine the efficiency of water use due to variations of weather. On the one hand, an 'optimistic' plant will produce a large leaf area, although in very dry years rainfall may be low or even zero so that the plant's survival is threatened. On the other hand, a 'pessimistic' plant ensures its survival and production. In general, an 'optimistic' behaviour, with high stomatal conductance values, is more productive, while a 'pessimistic' attitude with lower stomatal conductance values is more suitable in dry years (JONES 1983).

Other authors have studied seasonal water consumption or vine responses to soil water availability (BOTA *et al.* 2001, DELOIRE *et al.* 2004, MEDRANO *et al.* 2003). However, little is known about the vine water use after a period of water stress or the response of vines grown either under high or low water availability conditions to water stress. In this work we studied responses of two cultivars to watering previously being subjected to water stress. For this study two cultivars originating from different climatic zones were used: 'Airén' from the dry zone of La Mancha and 'Chardonnay' from the humid zone of Burgundy.

### Material and Methods

This experiment was carried out at the Universidad Politécnica de Madrid in Spain (40°26'36"N, 3°44'18"W, 590 m a.s.l.). Three-year-old grapevines were grown in 35 L lysimeters with a mixture of peat, sand and organic soil (63:25:12) and covered with a plastic film to eliminate evaporation and the infiltration of rainfall. The vines were grown open air in full sun. Drainage was measured in a second container. A 0.2 m TDR soil moisture probe (Trase Soil Moisture, Santa Barbara, CA, USA) was installed in each lysimeter. Each vine was restricted to two shoots. This experiment was completely randomised with 5 single vine repetitions. Two factors were analysed: cultivar and irrigation. The cultivars were 'Chardonnay' and 'Airén' grafted onto '1103 Paulsen'. Irrigation treatments started on May 8.

In the non stress irrigation treatment (NS) the potting medium was kept close to field capacity by applying throughout one week the amount of water the vines had consumed the previous week. Water consumption was determined gravimetrically by allowing drainage. In the stress treatment (S) we applied 50 % of the water consumed by NS vines for each cultivar with corrections for differences in leaf area between treatments, calculated by the following formula:

$$W_s = 0.5 \cdot W_{NS} \cdot LA_s \cdot LA_{NS}^{-1}.$$

Where  $W_s$  = water applied to stressed vines,  $W_{NS}$  = water applied to non-stressed vines,  $LA_s$  = leaf area of stressed

vines and  $LA_{NS}$  = leaf area of non-stressed vines. Leaf area per vine was measured biweekly. Leaf area was estimated by developing a second order polynomial equation, relating main vein length to leaf area of 30 leaves of each cultivar, as proposed by CARBONNEAU (1976).

On July 31 all experimental vines were watered to reach field capacity. No additional water was applied for the next 6 d. On August 1, 3 and 5, water consumption of 3 vines of each cultivar was evaluated in each irrigation treatment. Soil moisture content was measured on 5 vines of each cultivar and irrigation treatment. On July 26 and August 1 predawn leaf water potential ( $\Psi_{pd}$ ) of three vines of each cultivar was measured in both irrigation treatments. Before rewetting (July 26) and 1, 3 and 5 d after rewetting (August 1, 3 and 5) leaf conductance ( $g_l$ ), transpiration ( $E$ ), and net photosynthesis rate ( $A$ ) were measured at 11 a.m. (9:00 solar hour aprox.) using healthy, mature sun-exposed leaves. On July 26, August 1, 3 and 5 at 11 a.m. relative humidity was 24.5, 35.3, 32.4, 28.3% and temperature was 28.1, 28.7, 24.3, 24.8 °C. Predawn leaf water potential was measured with a pressure chamber (Soil Moisture Equipment, Santa Barbara, CA, USA). Transpiration ( $E$ ) and leaf conductance ( $g_l$ ) were measured using a steady state porometer (Li-1600, Li-COR, Lincoln, NE, USA). Net photosynthesis was measured with a portable IRGA system (Li 6200, Li-COR, Lincoln, NB, USA) on leaves fully exposed to solar radiation.

Variance analyses were carried out using the software package MSTAT-C (University of Michigan).

## Results and Discussion

**Water consumption:** One day after rewetting (August 1) the soil water content did not differ significantly between treatments or cultivars (Tab. 1). However, 3 and 5 d after rewetting soil water content was significantly higher in water-stressed vines than in non stressed vines and 'Chardonnay' had higher values than 'Airén'.

Table 1

Predawn leaf water potential ( $\Psi_{pd}$ ) the day after rewetting (August 1) and water content 1, 3 and 5 d after rewetting for 'Airén' (A) and 'Chardonnay' (C) grapevines cultivated without water stress (NS) and under water stress (S) before rewetting. Factorial analysis of variance (CV = cultivar, IT = irrigation treatment, CUL·IT = interaction)

	$\Psi_{pd}$ (MPa) August 1	Water content % (v/v)		
		August 1	August 3	August 5
C-NS	-0.20	25.3	9.5	7.5
A-NS	-0.17	25.1	9.5	4.3
C-S	-0.35	30.3	29.3	16.8
A-S	-0.25	26.9	21.3	10.2
CV	*	ns	*	**
IT	**	ns	**	**
CV·IT	ns	ns	*	ns

ns, \*\*, \*, non-significant, significant at  $P = 0.01$  and  $P = 0.05$ , respectively.

Water consumption during the 5 d after rewetting was directly related to vine leaf area (Fig. 1); no significant differences were found between cultivars or treatments (statistical analysis not shown). Unstressed 'Airén' vines ('Airén' NS), which developed a larger leaf area before rewetting, consumed more water than those that had developed a smaller leaf area ('Chardonnay' S). In the NS and S treatments leaf area development was greater in 'Airén' than in 'Chardonnay'. The leaf area of non-stressed vines was 3 and 2 times greater than that of stressed vines in 'Chardonnay' and 'Airén' respectively. In stressed vines leaf area developed from budburst to fruit set in both cultivars. In NS vines however, leaf area developed from budburst to veraison (GÓMEZ DEL CAMPO *et al.* 2002).




Fig. 1: Water consumption during 5 d after rewetting as a function of vine leaf area ( $Y = 2.78x + 2.68$ ,  $R^2 = 0.68^*$ ) for 'Airén' and 'Chardonnay' grapevines cultivated without water stress (NS) and under water stress (S) before rewetting.

**Predawn leaf water potential ( $\Psi_{pd}$ ):** Although the amount of water available to vines after rewetting was not significantly different, significant differences in  $\Psi_{pd}$  were found between treatments and cultivars the night after rewetting (Tab. 1). Significantly higher  $\Psi_{pd}$  values were obtained in NS vines than in S vines. 'Airén' showed significantly higher  $\Psi_{pd}$  values than 'Chardonnay' in both treatments. Higher  $\Psi_{pd}$  in NS indicate that, although soil water availability after rewetting was the same in both irrigation treatments (Tab. 1), vines subjected to water stress before rewetting are unable to reach the hydration levels of non-water stressed vines. Sap cavitation in xylem may have prevented total hydration of the water-stressed tissues, thereby causing differences between cultivars as was observed by SCHULTZ (2003). Moreover, the lower development of the root system of water-stressed vines (GÓMEZ DEL CAMPO *et al.* 2005) may explain the lower re-hydration capacity. Other authors have also observed that the recovery of vines after rewetting is not immediate. In vines cultivated in pots under water stress conditions,  $\Psi_{pd}$  recovered 2 d after rewetting (HARDIE and CONSIDINE 1976). In a field study SCHULTZ (1996) observed that  $\Psi_{pd}$  only slowly recovered after a prolonged period of drought and one precipitation event.

Cultivar differences in rehydration during the night have been shown by BOTA *et al.* (2001), MEDRANO *et al.* (2003) and GÓMEZ-DEL-CAMPO *et al.* (2004). SCHULTZ (1996) found that  $\Psi_{PD}$  of different cultivars responded differently to one precipitation event after a prolonged period of drought. SCHULTZ (2003) attributed these cultivar differences to differences in the hydraulic conductance of stems and particularly petioles. He obtained different cavitation rates between grapevine cultivars and reported that these differences were related to vine responses to water stress; cultivars with greater hydraulic conductance were more sensitive to cavitation which induced stomatal closure. Besides the hydraulic control of stomatal action there is strong evidence for a hormonal control, mainly by ABA, (CORREIA *et al.* 1995, LOVISOLO *et al.* 2002).

**Stomatal behaviour:** Before rewating  $g_i$ , E and A values of unstressed vines were three times greater than those of the S treatment (Tab. 2). After rewating leaf conductance increased in both irrigation treatments, 1.7 times in NS and 3.7 in S vines. The high value

Table 2

Leaf conductance, transpiration and net photosynthesis at 11 a. m. before rewating (July 26), 1, 3 and 5 d after rewating for 'Airen' (A) and 'Chardonnay' (C) grapevines cultivated without water stress (NS) and under water stress (S) before rewating. Factorial analysis of variance (CV= cultivar, IT = irrigation treatment, CUL·IT = interaction)

	July 26	August 1	August 3	August 5
Leaf conductance ( $\text{mmol H}_2\text{O}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ )				
C-NS	123.6	267.6	83.8	129.2
A-NS	179.4	229.2	92.1	108.7
C-S	51.0	180.2	149.8	134.7
A-S	37.5	146.4	167.5	163.2
CV	ns	ns	ns	ns
IT	**	**	ns	ns
CV-IT	*	ns	ns	ns
Transpiration ( $\text{mmol H}_2\text{O}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ )				
C-NS	4.03	7.08	1.72	3.37
A-NS	5.58	6.44	2.11	2.85
C-S	1.86	5.55	3.38	3.21
A-S	1.46	4.47	3.65	3.86
CV	ns	ns	ns	ns
IT	**	*	*	ns
CV-IT	*	ns	ns	ns
Net photosynthesis ( $\mu\text{mol CO}_2\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ )				
C-NS	11.8	15.4	9.8	8.7
A-NS	11.8	14.5	9.0	4.5
C-S	4.8	11.5	10.7	11.7
A-S	3.7	10.6	11.3	12.3
CV	ns	ns	ns	ns
IT	**	**	ns	*
CV-IT	ns	ns	ns	ns

ns, \*\*, \*, non-significant, significant at  $P = 0.01$  and  $P = 0.05$ , respectively.

in NS vines on August 1 compared to June 26 may be due to relative humidity which increased from 24.5 to 35.3 %. One day after rewating  $g_i$  was significantly higher in NS vines (Tab. 2). However, no significant differences were found between treatments in any of the forthcoming measurements. Three d after rewating a sharp decrease in  $g_i$  was observed in the NS vines, nevertheless S vines maintained similar  $g_i$  values during the 1, 3 and 5 d after rewating. This response of  $g_i$  in NS vines to a punctual rewating followed by 6 d without irrigation was related to the high water consumption between August 1 and 3 (Tab. 1) due to the high leaf area which had developed due to the previous water availability (Fig. 1). Stressed vines had more stable  $g_i$  values after rewating (Tab. 2). During the 6 d without irrigation,  $g_i$  values were not as low as those of S vines before rewating (on July 26). At 11 a.m. E values were significantly different between treatments at the first two measurements after rewating (Tab. 2). One day after rewating the NS treatment had significantly higher E values, while the S treatment had significantly higher values 3 d after rewating. One and 5 d after rewating A values were different between treatments (Tab. 2). One day after rewating unstressed vines had significantly higher values, while these values were significantly lower 5 d after rewating. Other authors have also observed that the recovery of  $g_i$ , E and A after rewating of stressed vines was not immediate (KIEWER *et al.* 1985, LOVEYS and KRIEDEMANN 1973, PONI *et al.* 1993), although these values recovered more rapidly than  $\Psi_{PD}$  (SCHULTZ 1996). Previous to rewating S vines were subject to severe stress, after rewating, the experimental vines were subject to mild stress considering that  $g_i$  was higher than or close to  $150 \text{ mmol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$  (CIFRE *et al.* 2005).

$g_i$ , E and A were not significantly different between cultivars during the experiment (Tab. 2).

Therefore, the lower hydration of vines subjected to water stress before rewating (Tab. 1) allowed these vines to maintain constant  $g_i$  and A rates after rehydration as well as significantly higher A rates 5 d after rewating (Tab. 2).

**Relationship between  $g_i$  and A:** Significant differences in the relationship between  $g_i$  and A were found between water treatments before rewating (Fig. 2). Related to  $g_i$ , A values were higher in the NS treatment, which confirms results of CHAVES (1986), BOTA *et al.* (2001), GÓMEZ-DEL-CAMPO *et al.* (2004). After rewating however the relationship between A and  $g_i$  was not significantly affected by previous water conditions (Fig. 3). This seems to indicate that the relationship between A and  $g_i$  is dependent on the water availability. Differences in the adjustment of A to  $g_i$  seem to be a specific response to a specific situation of water availability and should not be considered a permanent adaptation mechanism.

After rewating, the water use efficiency ( $A/g_i$ ) was not significantly different between treatments or cultivars (Tab. 3). This confirms that it is preferable to use the term 'drought tolerant' rather than 'drought resistant' to describe the mechanisms allowing plant survival or productivity under drought conditions (JONES, 1983).




Fig. 2: Net photosynthesis as a function of leaf conductance at 11 a.m. before rewatering of 'Airén' and 'Chardonnay' grapevines cultivated without water stress (NS) and under water stress (S) before rewatering. NS:  $Y = 1.9 \ln(x) + 2.2$   $R^2 = 0.15$ ; S:  $Y = 1.7 \ln(x) + 1.6$   $R^2 = 0.60^{**}$ .




Fig. 3: Net photosynthesis as a function of leaf conductance at 11 a.m. 1, 3 and 5 d after rewatering of 'Airén' and 'Chardonnay' grapevines cultivated without water stress (NS) and under water stress (S) before rewatering. NS:  $Y = 4.3 \ln(x) - 9.0$   $R^2 = 0.90^{**}$ ; S:  $Y = 4.4 \ln(x) - 9.7$   $R^2 = 0.58^{**}$ .

### Conclusions

The only permanent adaptation mechanism to water stress observed in this assay was the lower leaf area development in S vines which allowed to use less water and to maintain higher water availability levels than NS vines. After rewatering S vines did not rehydrate like NS vines possibly due to cavitation and/or lower root development. As a result of lower leaf area and lower hydration, stomatal conductance in S vines remained fairly constant and 5 d after rewatering A was higher than in NS vines.

The use of  $\Psi_{pd}$  as an indicator for grapevine irrigation management proposed by CARBONNEAU (1998) should not only consider phenological stages (DELOIRE *et al.* 2004) and cultivars (BOTA *et al.* 2001, MEDRANO *et al.* 2003, SCHULTZ

Table 3

Water use efficiency (A/g<sub>l</sub>) at 11 a.m. 1, 3 and 5 d after rewatering on July 31 for 'Airén' (A) and 'Chardonnay' (C) grapevines cultivated without water stress (NS) and under water stress (S) before rewatering. Factorial analysis of variance (CV= cultivar, IT = irrigation treatment, CUL·IT = interaction)

	A/g ( $\mu\text{mol CO}_2 \cdot \text{m}^{-2} \cdot \text{s}^{-1} / \text{mmol H}_2\text{O} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ )		
	August 1	August 3	August 5
C-NS	0.058	0.168	0.135
A-NS	0.064	0.332	0.010
C-S	0.072	0.081	0.300
A-S	0.073	0.071	0.076
CV	ns	ns	ns
IT	ns	ns	ns
CV·IT	ns	ns	ns

ns, non-significant at  $P = 0.05$ .

2003, GÓMEZ-DEL-CAMPO *et al.* 2004), but also previous water conditions of the vine.

In spite of the significant differences in  $\Psi_{pd}$  found between cultivars after rewatering,  $g_l$ , E and A values were found not to be cultivar specific. Due to its stomatal behaviour under conditions of prolonged water stress, 'Chardonnay' can be considered 'luxurious' while 'Airén' could be considered 'pessimistic' (GÓMEZ-DEL-CAMPO *et al.* 2004) according to the classification of BOTA *et al.* (2001), or they could be considered 'isohydric' and 'anisohydric', respectively, terms suggested by SCHULTZ (2003). These genetic differences in stomatal behaviour however, disappeared after rewatering. Likewise, the relationship between A and  $g_l$  was not significantly different between treatments after rewatering.

Vines subjected to high water availability experience a sharp decrease in A when irrigation is stopped due to the high water consumption by the great leaf area and high hydration capacity. These results could help to develop irrigation scheduling under conditions of water shortage. Long irrigation intervals with application of high amount of water will cause lower levels of A when vines have consumed the available water than applying low amount of water frequently, because it will induce adaptation of the leaf area development to the available water, thereby improving water use efficiency.

### References

- BOTA, J.; FLEXAS, J.; MEDRANO, H.; 2001: Genetic variability of photosynthesis and water use in Balearic grapevine cultivars. *Ann. Appl. Biol.* **138**, 353-361.
- CARBONNEAU, A.; 1976: Principes et méthodes de mesure de la surface foliaire. Essai de caractérisation des types de feuilles dans le genre *Vitis*. *Ann. Amélior. Plantes* **26**, 2, 327-343.
- CARBONNEAU, A.; 1998: Irrigation, vignoble et produits de la vigne. In: J. R. TIERCELIN (Ed.): *Traité de Irrigation*, 25-37. Tec & Doc, Paris.
- CHAVES, M. M.; 1986: Fotossíntese e repartição dos produtos de assimilação em *Vitis vinifera* L. Doct. Diss. Univ. Lisbon, Portugal.



- CIFRE, J.; BOTA, J.; ESCALONA, J. M.; MEDRANO, H.; FLEXAS, J.; 2005: Physiological tools for irrigation scheduling in grapevines (*Vitis vinifera* L.). An open gate to improve water-use efficiency? Agric. Ecosyst. Environ. **106**, 159-170.
- CORREIA, M. J.; PEREIRA, J. S.; CHAVES, M. M.; RODRÍGUEZ, M. L.; PACHECO, C. A.; 1995: ABA xylem concentrations determine maximum daily leaf conductance of field-grown *Vitis vinifera* L. plants. Plant, Cell Environ. **18**, 511-521.
- DELOIRE, A.; CARBONNEAU, A.; WANG, Z.; OJEDA, H.; 2004: Vine and water: A short review. J. Int. Sci. Vigne Vin, **38**, 1-13.
- GÓMEZ-DEL-CAMPO, M.; RUIZ, C.; LISSARRAGUE, J. R.; 2002: The effect of water stress on leaf area development, photosynthesis and productivity in Chardonnay and Airén grapevines. Am. J. Vitic. Enol. Vitic. **53**, 138-143.
- GÓMEZ-DEL-CAMPO, M.; BAEZA, P.; RUIZ, C.; LISSARRAGUE, J. R.; 2004: Water stress induced physiological changes in leaves of four container-grown grapevine cultivars (*Vitis vinifera* L.). Vitis **46**, 99-105.
- GÓMEZ-DEL-CAMPO, M.; BAEZA, P.; RUIZ, C.; LISSARRAGUE, J. R.; 2005: Effects of water stress on dry matter content and partitioning in four grapevine cultivars (*Vitis vinifera* L.). J. Int. Sci. Vigne Vine **39**, 1-10.
- HARDIE, W. J.; CONSIDINE, J. A.; 1976: Response of grapes to water-deficit stress in particular stages of development. Am. J. Enol. Vitic. **27**, 55-61.
- JONES, H. G.; 1983: Plants and Microclimate: A Quantitative Approach to Environmental Plant Physiology. Cambridge Univ. Press., U.K.
- KLIEWER, W. M.; KOBRIER, J. M.; LIRA, R. H.; LAGIER, S. T.; DI COLLALTO, G.; 1985: Performance of grapevines under wind and water stress conditions. In: D. A. HEATHERBELL; P. B. LOMBARD; F. W. BODYFELT (Eds): International Symposium on Cool Climate Viticulture and Enology, 198-216. Oregon, USA
- LOVEYS, B. R.; KRIEDEMANN, P. E.; 1973: Rapid changes in abscisic acid-like inhibitor following alterations in vine leaf potential. Physiol. Plant **28**, 476-479.
- LOVISOLO, C.; HARTUNG, W.; SCHUBERT, A.; 2002: Whole-plant hydraulic conductance and root-to-shoot flow of abscisic acid are independently affected by water stress in grapevines. Funct. Plant Biol. **29**, 1349-1356.
- MEDRANO, H.; ESCALONA, J. M.; CIFRE, J.; BOTA, J.; FLEXAS, J.; 2003: A ten-year study on the physiology of two Spanish grapevine cultivars under field conditions: Effects of water availability from leaf photosynthesis to grape yield and quality. Funct. Plant Biol. **30**, 607-619.
- PONI, S.; LAKSO, A. N.; TURNER, J. R.; MELIOUS, R. E.; 1993: The effects of pre-and post-veraison water stress on growth and physiology of potted Pinot Noir grapevines at varying crop levels. Vitis **32**, 207-214.
- SCHULTZ, H. R.; 1996: Water relations and photosynthetic responses of two grapevine cultivars of different geographical origin during water stress. Acta Hort. **427**, 251-266.
- SCHULTZ, H. R.; 2003: Differences in hydraulic architecture account for near-isohydric and anisohydric behaviour of two field-grown *Vitis vinifera* L. cultivars during drought. Plant, Cell Environ. **26**, 1393-1405.

Received June 13, 2006

